

HOME CARE FOLLOWING ENTROPION/ECTROPION REPAIR, EXCISION

OF LID LESION WITH SKIN GRAFT

POST-OPERATIVE CARE

The following is a list of instructions for you to follow now that you have had your surgery. This will ensure the healing process goes smoothly and no complications arise

Expected Pain and Management

 When your anaesthetic wears off, you may feel some discomfort in your eye; take Panadol or Pandeine to relieve this.

Dressings

- You may take the crepe bandage off four (4) hours after discharge.
- The eye dressing <u>MUST NOT</u> be removed until reviewed by your doctor 5-6 days postop.
- Leave other dressings in place until your review appointment.
- It is important not to rub the eye.

Daily Care:

Some non dissolvable stitches will be removed at your first and subsequent post op reviews. Other stitches WILL DISSOLVE. They do not need to be removed.

Wound Care

Once the dressings are removed by your doctor, if there is any discharge, dab the
margin of the lids with a clean tissue or cotton bud dipped in warm, boiled water.
Discard tissue after each wipe.

General Care:

- Ensure you have adequate rest until you feel capable of resuming normal activities.
- Resume your normal diet as soon as you feel inclined to do so.
- Avoid alcohol 24 hours following an anaesthetic.
- Shower or bathe as usual, keeping water out of the eye.

NO vigorous activity for 2 weeks.

It is normal for the eye to be uncomfortable for the first 4 weeks – normal pain relief (e.g. Panadol/Panadeine or Nurofen) is fine to take if needed.

Contact your doctor if any of the following occur -

- Unrelieved pain/swelling in the eye.
- Excessive discharge requiring more than two (2) tissue wipes per hour after 1st post op day.

If you have any concerns, you can ring anytime for an earlier appointment.

Emergency Phone Numbers		
South West Eye Surgery	5562 4488	
South West Health Care	5563 1666	
Royal Victorian Eye & Ear Hospital	99 298 666	
FOLLOW UP APPOINTMENT – 1 week		
Doctor		Date
Place		Time